



UNIVERSITY OF CALIFORNIA MUSEUM OF VERTEBRATE ZOOLOGY

Vol. 262










Ogilvie, F. 1942 Kuigs Canyon Natral Park, Culif.



Catalog of Specimens



17.

Cedar Grove, South Fork Kings River, 4600 ft., Fresno Co., California may 21, 1942 & Deromyscus maniculatus 131-58.5-20-15, may 22, 1942 6 Geromyseus maniculatus 150-65-21-19, 19.2 oms & Thomomys bottse 212-22-67-8, 118.3 óms 5 Sceloporus graciosus 3 Sceloporus graciosus may 23, 1942 6. 9 Deromipeus maniculatus 158-68-20-20, Sceloporus graciosus May 24, 1942 8 9 Geromipeus maniculatus 190-95-23-21, 27.3 gms. Sceloporus graciosus May 25,1942 10. 8 Peromyseus maniculatus 157-71-20-20, 23.2 gms Near Ceder Grove on trail to Zumwalt Meadow, 4600-5000 ft. Fresno Co. Calif. 11. Sceloporus graciessus 12 6 Sceloporce 13. 8 Sceloporus Zum beelt Meadows, Rocks immed. below brand Sentine, 5000 ft, Fresno Co; 4 Sceloporus Zumwaet Meadows, 5000 ft., com how how here to, Calif.

. if me

Catalogue
May 30,1942

Cedar Grove

discarded

discarded

18	1/4 Marie Ma	
	Sceloporus undulatus	
	9 Scelspour graciosus	
21	8 Scelopous graciosus	
	& Sceloporus graciosus	
	3 Suloporus graciosus	
24	8 Sulapous graciosus	
	8 Sceloporus graceosis	
26	3 Sceloporus graciosus	
	4 Scelopous graciosus	
	Sceloporus graciosus	
妈	May 31,1942	2006
29		20.6
30	8 Beromyous maniculatus 130-52-20-18	10.8
<b>1</b>	3 Geromyseus maniculatus 116-52-19-17	7.6
32		
<i>3</i> 3	13 mi. " - " - "	



## Journal

May 21

Cedar Grove, South Forktings R., 4600 st., Freano Co., Calif. Picket up 15 traps set the night before at Ta.m. They were located among dead logs on south side of lings Sivar Whole region was marshy with areas of dry land. Only one Berympeus maniculatus was found a young specimen (1). Those traps which contained nothing appeared not to have been touched. The collected specimen was prepared. about 1:15 p.m. Mr. Disowled the group to Sheepikun Falls. On the way up many yellow Pines and misery plant were seen. About 4 Sceloporus were seen scampering very actively on dead logo. None were caught, but a small 2) specimen was, it was Sceloporus graciosus. Otale Falls (5200 ft.) the nest of a Caliope Hummingbird was seen in pinetree (with a eggd. The female was seen by some members of the group, but was not present when seen by me. Continued up along wellmarked trail. Mt. Sardner could be seen clearly; it was covered with snow. Mt, Fordner is app. a good adays hike from Cedar Grove. The crest of the Sierras is about the same distance beyond Mt. Gardner (Dison). Two White Throat Swifts were seen oflying above. Their long wings made them very distinct. Yellow Pine & Sugger Bine were the most abundant trees. There were some Manganita and much Misery Plans, Some Blackthroated Grays were heard. Further up a Junco was seen. It's head was much blacker of body much lighter than the Berkeley race. Its song was peculiar and not very characteristic, this was noticed ly Dr. Wille. achipment was seen in togdead trunk - head was protruding from hole + watching us. Blids seen were: Nedtailed Hawk, Calveras Warbler, Ecleolated Warbler, Elack-throated growy warbler, Stellas Jays, Brown Creeper, White throatel Swifts, Mt. Chickadee, Cassin Turple Finch, White headed Woodpecha, Caliope Humminglind. Many of these were not seen too clearly by me.




may 22

Cedar Grove, 3. Fork Kings R. Hooff,
Collected the 11thops at 6:15 that had been set the night before
up on dry flat just below beginning of hills to west Jeamp (400 yg).
Nothing was found, but 2 traps had been spring of the bait
gone. One trap had lost the bait but was not spring.
Loter in morning accompanied H. F. T. D. de M. to set gopher traps.
6 were set by digging between the mounds of placing double
traps

Trocks

In order to trap gopher


coming in either direction. Went back about 2 hrs. later (11: ooah and found 2 gophers, trought them back & d prepared the male (3). It was found on n. side of King VI. in drag gravely soil, amid yellow pines, misery plant, etc. There were many old fallen trees about. In late afternoon (4:00) climbed small hill to west of camp in search of ligards. among grainte outeropping a number of Sceloporus were seen scampering about. There were fallen pines near by white with weathering; lizards were also seen about these. A skink was seen in the crevice of a large rock, but it slipped thru the noose. Two Sceloporus (4+5) were finally caught. Continuing north east a few hundred yards we came upon a rattlesnake tightly coiled in a dead log. It remained perystill & did not ratile although a stick was throwned the log. Finally the range Mr. Gimm, was trought & he Killed the rattler Ten trops were set: 9 by the west side of the bridge among the grass of old logs, I was set about 100 yels from lang



May 23

Journal

Cedar Grave S. Fork Kings R., 4600ft., Fresno Co., Calif. at 6:15 collected the 11 trops set night before. none I those down by the river had been snapped. In a trop nearly, set by another, a microtus & was found. It was lost from the pocket of the collector & could not be found. In a trap set about 150 yds. from camp under an old dead pine steemp a young of Gerompous was found. about 2:30 went to same group of rocks a short distance up hill to west of camp. after several minutes only 1 Sceloporus was seen whereas the day before there had been 4 or 5. Went down to same dead log where the rattlesnake had been found. A beautiful skink was seen, but it got away. Also I saw a Schoporus which was practically black, it also escaped. Then went about a hundred or so yards up Sheep breek tail and caught one Sceloporus graciosus 3 (7) The reattlesnake was skinned and the unner surface rubbed with alumo arrenic. It was nailed in the shade on a pinetree & left to dry. The snohe was dissected. It was 91 km. long.



## Journal

may 24

Cedar Grove, S. Fork Kings N., 4600 ft., Fresno Co., Calif. a q Gerompeus was collected by trunk for other side of bridge. The other traps which were set along the Vack Station road along dead lop, etc. yelded nothing. The back was same or not tampered with but the top of a Black Oak a Fanager was observed singing. Below on same limba warbling Vired. Both were watched for some mins. In afternoon what in search of lizards. The skink by the granite out cropping was seen and many attempts were made to eatch it. It was a tight dark brown with yellowish stripes down back Jail was also dockish in color. It was fairly alow moving but it had such a narrow head o body, I was slippery exough to slip out of the moose. another skink with a blue tail was seen in burndlog. While in this region a large coyste - tan to huff color with a large bushy tail tipped with black - came trotting by. It was seen again a short distance from this area. The ranger later said it was a frequent visitor to the park. at foot for the Creek tail a youngish Sceloporus was caught. at 5:00 pm. set about 12 traps baited withmest, downly extrance of Sheep Creek into King K and along the grassy meadow some yes from the King T. to the southeast of the camp. Mr. + Mrs. Dixon ate supper with the group. 6 trout had been donated by neighboring campers. Jean Russich & I then set 31 traps baited with nuts or oatmest peanut butter mix down by the ranger's harn and neighboring region. On way back wend to the Robin's nodon for side of bailge accidently scared of Tobin from nest. We watched some distance away. She flew to nearly pine and come a steady, single call note. after a while This note subsided she began & presn herse. & arrived after about 15 min. In 30 min. neither had gotton on nest.



Journal May 25 Cedar Geore, S. Fork King R., 46 ouft., Freeno Co., California Only one Serompeus was caught at the trown. None of the traps by the nove downiesled anything, although 31 traps had been set. It was noticed that 30,74 garbaje cans had been turned over & contents messedahout; later apparently fresh bear scat was found about 20-30 yds. from eamp. Nogefinite foot fruits were seen. at 1:00 Mr. Dison led us to Zumwalt Meadows about 5 miles from camp. Cach person had approximately 30 mus. sp. traps 43 rattraps. be drove car about 2 mi. and then hiked rest I way Fail was fairly level there was a very gradual clink gahout 500 ft. as zumwalt Mealows is around 5000 ft. Heard several Chipping Sparrows hud did not see them did see trus white headed boodpeckers very distinctly. The close were climbing up trunk high up ina pine tree; they were close together. a cassin Vires was head singing ingroved pine ocedar. Laty on waytome one was seen closely. It was medium, graycolor. along truit nany yellow fines & some su jaspines were seen. Also several sinon pines were seen. Two Calif. mule deer does wen seen in thepines. The large nests of 8 or 9 woodrats were observed. Induay home 8 traps were set - scattered by several of these. a & Blackthroated Stay worden was seen on a low link of a pine; This was in a densely wooded area, also farther along path a Toursend Solitaire was seen along trank of the river It had a long narrow neck thead, trim lines, o greijsh trown color. When it flew white stripes on the wings could be seen. Zumwalt medowe is a large owampy area, thickly covered with tallgrass, & cot tails. Much of the water appeared to be stagnant. Along banks ever many willows & dogwoods. Grand Sentine Mt. Tuerlooked the meadows. Boulders which had fallendown from Dr. Sentinel lined one side of the meadow closely. Several Sceloporus undulatus were caught here. 26 traps were.



Journal

May 25

Cedar Grove, S. Fork Kings R., 7600ft., Fresno Co., Colif set along south side of the meadows. A number of Redbing Blockbuds were nesting in meadows. The 3" were conspecuous on Willows, etc. around. I's were occasionally seen. They are brownish with brown or white speckled breast. Three or four bline- green swallows were flying across modow. Song spacrows were present also. They have a lovely song different from Backeley Syranows. a allen Humminghird was heard of then seen on very top twig ga willow tree. On way back several ligards were caught, and hirds of same type as those seen coming to meadows. It was clouded & rained slightly on way home. I kat traps were set by woodrat houses.

Left camp at 9:00 am to check traps. Mr. Dixon led way. He Killed a Hyratcher with a police revolver using dust shot. The stuffed mouth with cotton immediately to prevent digestive fluids from running out & soiling feathers. He then placed bird head downward in come of paper V. Clashe Killed a peleololad harbler 3-bright yellow treast, black top of head. Sawpeverd wood Gewees (flycation). Nests are usually found in fork of tree. many service berries were in bloom. Whentrops were reached, nothergross found in mine, However a Murstus and a Teromysus truei suere caught. Mr. Newon shot a song Sparrow. I caught a Sceloporus ungulatus on the large rocks below Grand Sentine, across river a harge rubber make was caught, white in the meadow a allegator lizard was caught.



Ogilvie 1942

## Journal

may 27

Cedar Grove, S. Fork TingoR., 46 ooft, Gresno Co., Calif.

Left camp 25 7. ovam wich Mr. Dijon for zumward mes down Daywak clear toold. When we passed by kattraps by woodrat hole, nothingwar found. Saw 2 heartful thay squirtrels in tall pine tree. Beached zumwalt of found none of my trops spring. One member found a Larterstake Mr. Dijon shot a bory sparrow, a Q Redwingel Blackbird, a Calif. Linnet (Gurple Finch), & a western flycother. I attempted to catch a large & Sceloporus ungulatus on the rock behind meadow, but 2 god away. Beturned home for lunch 1:00 pm. ahout 6:00 pm. Mr. Dijon led group & mt. side to northcast yearsp. Fraps were set as follows—lack person keeping a district trap line:

I set 15 trup in mo. 2 position - placing them among thuck bushes & beside rocks.

May 28

Left campat 7: to am Bochick traps on hill. Ched & clear morning but low clouds on nountains. Found that none of my traps had been touched. Left them as they were. On way home saw a water legal (Dipper) fly from under bridge. Came hack a sat under bridge. Saw the Dippers nest under girders. It q were standing on rocks bobbing up & down. It q were standing on rocks bobbing up & down. It windshop would grey wich very short tacks. Later male was seen feeding female. The flew to nest & entered from side entrance. Jean Aussick & I hiked at 9:00 am downeast side of river. Ate lunch on way. Hited past Zumwaed Meadows.



Cedar Grane, S. Fork Ringo R., 7600 ft., Freeno Co., California ded not see much animal life on trail. One grey squire was seen closely - eating a mut of some sort. also saw a 844 Chipping May 29 Left campat 7:45 with Mr. Dison. Headed up old trail to

Summit Meadows, 8000 ft., then liked to Kookout Seate, 8,470ft, and ate lunch on the grante peak. Teturned by following deer trails down varound to other side of mountain. Walked thru prow fields & took several pictures of the region. Came upon the new trail and followed it to Sheeps Creek. Watched & Calliofe Humminghird on nest for about 15 min. Seturned to compations 4: 30. We had hiked approximately 4 miles, climbing about 4000 ft. On this hike several new hirds & plants were seen. Sheer Auttal Hycatcher, sichowas heard first for some some before seen, had a song - . also a Sierra Creepes was seem hopping up trunk of a clead pine. a tay Sparrow was also seen & heard often & clearly. Bird is reddish brown above - breast is whitish with triangular brown spots. also the Vilolatel Woodpecker was heard hub not seen by me. The mt, Quail call was heard several times. Cet Summit meadows the Rosetreasted Sapsricker was seen on a dead Lodgepole Vine. 7 dyla (?) was also presend in the meadows. Many Robins were also seen heard. While on the top of Mit. Lookaut the Clarke Nuteracker (or crow) was seen - color were vrey, black, & white Severolnew plants were seen - Jeffery Vine, Snow Dush, Lodgepole Pine, & White veolets.



May 29

Species seen: Fox Sparrow
Western Bohin Gileolated Woodpecker
Western Janoger Mr. Chickodee
Coloners Warbler Mr. Quail (heard)
Dregon Junes Gose breated Sopsucker
White Arosted Swift Black headed Dorbesh
Muttall Sharolcher Clarke Nutcracker
Sierra Creeper Callished winninghird
Checked the 15 traps on hielside which had been

may 30

Checked the 15 traps on hillaide which had been set May 27.


3 of them had been snapped, but no animals were caught.

Lover in morning took taillunch and with B. E. Pauleon went toward Jumuset Mealows. Caught 9 Sceloporus graciosus on the way. We ate lunch on rock on far side of Braining River. While climbing to top of this hill a Sceloporus undulatus was caught. Further down trailan alligator lijerd was caught. Several Brey Squirrel were seen on route. Returned & campalisant 5:00 p.m. After supper 14 traps were set on far side of river 4 across road. A coral king Snals was seen on road.

24 7:30 am found that of the traps on this sile? road 3 had been snapped with the bait gone. On far, side 3 road by a forther prosent young. Perongseus was found. It had

May 31

by a follow from they? Teromysens was found. It had been caught verose a leg & his tail had bleed. Several feet from the rock by a sld log an adult of was caught. Farther away from roads across small stream another young of Beromyseus was caught. The 3 were brought back to camp of put up. Hen set a line of trapsup hiel to back



May 31

Cedar Grove, S. Fork Kings R., 4600 ft., Freno Co., Calif.

If comp ( at 5:00 p.m. walked up to Sheep Creek Falls +

observed Calliope Humminghinds. & was disturbed by our

presence & flew about \*\* from branch to branch I

several trees. Ofter supper trap lines were inspected.


Several had been snapped but nothing caught. Teaplines

were left.


Journal

June 1


Traplines were followed and 3 young Geromacus were found in line on other side of river. Nothing was found in traps set on hilling back of camp. Peromyceus were given away as there was not time to prepare skins. B. E. Paulson T I left camp about 9:30 am. 7 mi. along highway (4000 ft) a Yellow racer was found dead in road. 13 miles along highway a Coral King Snahe was likewise found. Ate lunch at Omar Khayham in heins. Saw 4 Calif. Redreaded woodpeckers on fence by road just a few miles out of Fresno. Itophelat drugstore in Fresno & hought some rubbing alcohol for snakes. Left Fresno about 2:00 p.m. & drove thru Josemite Nat. Fark. Arrived in Berkeley about 1:30 cm. hy way of Merced, Calif.



Species accounts



Ogilvie 1942 Thomomys bottal May 22 Cedar Grove, S. Fork King R., 4600 ft., Fresno Co., Calif. a & Thomomys was collected by a gopher trap placed incenter of runway. The mound was on north side of Kings R. amid old fallen logs; soil was light colve, coarse, & dry. There was some grass and misery plant about. The trap was set about 9:30 am. 8 the Thomomy was collected between 11 + 11:30. There were cut stems in his pockets - all were approximately came size



## Crotslus

May 22

Cedar Grove, S. Fork Kings River, 74600 ft., Teono Co., Calif. 3:20 p.m. a rattlesnake was found coiled in the crevice of an old fallen pine which had appearance of being burnt. Snake was corled in atleast two coils (9); tail was not visible and no rattling was heard. a stick was thrown against the log. Inake stuck tongue out quickly several times of then held it out for a longer period vibrating it. Color of snake is light tan background with ( ) dark brown spots down back. Gale yellowish splotcher are irregularly placed about dark brown ones. Eyes are yellow. Fallen pine is about 50 yds. from paved road; other yellowpenes are growing about. Manzinita bush and very young Incense cedare are about. Ground is day a granely. We got very close to him but he held completely still. Did not even stick out tongue (did so only when stick was thrown of log.) The Langer, Mr. Dimm, came to kill the rattler. It had uncoiled itself but was in the same crevice of the long. a large stick waspoked at it of it for the first time began to rattle rattling continuous and crawl in straight path down middle of log. It was hit at again, othis time it crawled upinto a bush which was hanging over an end of the log. Over a half of its body would be held upright from a branch, mit a smaller portion holding to branch. It was knocked down to ground. I struck at stick - a small piece I the back from the stick stuck on his fang as he drew away. The next blow landed on his head & tilled him. His rattlers were lost in the scuffle after descrition it was found to be a male The

may 23

testis were large. The upper part of the lungs were very reed-but no hemospage had occurred. The tracker was





Sceloporus graciosus

may 22 Cedar Grove, South Fork Kings R., 4600 ft., Tresnolo., Calif.

In the hotofternoon (about 4:05 (wastim)) went about 300 yds west of camp, climbed only a few yds. uphill to a flat dry area. Alound one large outcropping of granite several Scelupous were seen. They dashed quickly about the rocks or scurried from base of rocks over to a hear by dead log. Several sizes were seen. The area was very dry and many dead trees were lying about. Misery plant was abundant. Two Scelopous graciosus were noosed (4 +5).

may 23

a Sceloporus was noosed shout a 100 yds. uptrail to Sheep Creek Falls. It was on ampering over a weathered log and jumped to a granite rock. While on rock it lifted itself up & down by front legs about 10 times.

May 24

a & Sceloporus Was caught in an old log about 150 yds. from Can

•

Ogilvie 1942

## Cerompeus maniculatur

may 21

Cedar Grove, South Fork Kingott., 4600ft., Freeno Co., Calif.

One Peromipeus maniculatur was found in trap by dead log about 50 ft. from Kings R. arjoung specimen, graijish above - wt. 9.2 gms. Irop was set about 7:30 pm the night before.

may 22

another Peromyseus (2) was collected on morth side of Kings R. close by a dead log in march grass area. Hwas collected by Virginia D. Miller.

may 23

AF Geromipeus maniculatus was collected about 100 yds. from camp by an old pinetrunk. Area around had only a few pines; ground was sparsely covered with thin grass and small low plants; it was mostly covered with pine needles. This q had 3 pr. of well developed mammae. The vagina was large and very pink. A little blood was oo sing from the urinary popillaes. The trap had caught well up on the back and not in the pelvic region. When it was dissected the horns of the uterus each contained four distinct swelling, but the embryos were so minute as to be undistinguishable.

May 24

a q Gerompeus was collected several hundred yds. on north side of bridge. It was caught by an old upright pine trunk. Four very slight swellings in each horn of uterus.

May 25

May 31

At was caught at the barn behind Ranger's house. Three Geromepeus maniculatus, 2 very young ones & a

mature o. There were no embryos in the o



